

The Second Painlevé Equation in the Large-Parameter Limit I: Local Asymptotic Analysis

Nalini Joshi*

Department of Pure Mathematics

University of Adelaide

Adelaide Australia 5005

`njoshi@maths.adelaide.edu.au`

February 9, 2008

Short Title: Large-parameter Local Asymptotics of P_{II}

Abstract

In this paper, we find all possible asymptotic behaviours of the solutions of the second Painlevé equation $y'' = 2y^3 + xy + \alpha$ as the parameter $\alpha \rightarrow \infty$ in the local region $x \ll \alpha^{2/3}$. We prove that these are asymptotic behaviours by finding explicit error bounds. Moreover, we show that they are connected and complete in the sense that they correspond to all possible values of initial data given at a point in the local region.

1 Introduction

The asymptotic analysis of the six nonlinear second-order ODEs (ordinary differential equations) known as the Painlevé equations [1] has been a subject of both classical [2, 3] and modern [4, 5] interest. More recent interest was

*Research supported by the Australian Research Council.

revived by their importance in physical applications, particularly in statistical mechanics (see [6]), and by their close relationship to soliton equations (see [7, 8]).

The solutions of the Painlevé equations are, in general, highly transcendental functions known as the Painlevé transcendents. An advantage of asymptotic description is that they give explicit behaviours of such transcendental solutions in terms of known classical functions. Such descriptions also give insight into the asymptotic behaviours of the solutions of soliton equations [9, 10].

In this paper, we focus on the asymptotic analysis of the second Painlevé equation (or P_{II})

$$y'' = 2y^3 + xy + \alpha \quad (1.1)$$

in the limit as $\alpha \rightarrow \infty$. P_{II} is well known to be a symmetry reduction of the modified Korteweg-deVries equation (and other soliton equations, see e.g. [8]). In such a reduction, the parameter α arises as a constant of integration. Therefore, a large value of α is related to large values of initial data for the soliton equation.

Although there have been many studies [2, 3, 10, 11, 12, 5, 13, 4, 14, 15] of the Painlevé equations in the limit as their independent variable approaches a singularity of the equation, the asymptotic limit as a *parameter* approaches infinity has not been widely studied, with two exceptions. Kawai, Takei, *et al* [16, 17, 18] have studied the Painlevé equations extensively in such limits through their associated linear equations, called isomonodromy problems [19, 20]. Kitaev [21] has studied the double asymptotic limit of the second Painlevé equation as both $x \rightarrow \infty$ and $\alpha \rightarrow \infty$, again through an isomonodromy problem. However, both studies appear to be restricted to formal special solutions.

In this paper, we prove that in a region bounded by $x \ll \alpha^{2/3}$ the local asymptotic behaviour of the general solution as $\alpha \rightarrow \infty$ is given by elliptic functions. We also show that there exist special solutions whose local asymptotic behaviours are given by degenerate elliptic functions, i.e. singly periodic functions. These results are analogous to the results found by Boutroux [2, 3] for the Painlevé equations in the limit as $x \rightarrow \infty$.

However, the domains of validity are quite different from those that are known for the limit $x \rightarrow \infty$. In our case, these domains are full disks (punctured at poles and stationary points) of large radius, unlike the case $x \rightarrow \infty$ in which such domains are either bounded local patches or sectors of angular

width strictly less than 2π .

Our methods are direct and do not use the relation of P_{II} to an isomonodromy problem. Part of our motivation is to show that rigorous general asymptotic results can be obtained through natural direct methods without going through associated linear problems.

However, we do rely on a characteristic property, i.e. the *Painlevé property*, of the Painlevé equations to obtain sharp error bounds. This is the property that all movable singularities of all solutions of each Painlevé equation are poles. In particular, we use the direct proof of the Painlevé property of the Painlevé equations given in [22], to define local paths of integration and to estimate w on such paths. Error bounds can be obtained without assuming this property (see Boutroux[2]). However, its usage allows us to obtain improved error estimates.

The Painlevé equations also possess many other deep properties. For example, their solutions can be written in terms of entire functions [23], they possess exact special solutions in terms of classical special functions and transformations relating solutions to other solutions (see [8] for references). Asymptotics is a natural part of their study as nonlinear special functions.

1.1 Asymptotic Regions

The study of P_{II} for $\alpha \rightarrow \infty$ necessarily decomposes into three parts corresponding to three regions. First, consider a transformation of P_{II} given by a dominant-balance analysis as $\alpha \rightarrow \infty$, namely

$$\begin{aligned} y(x) &= \alpha^{1/3} w(z) \\ z &= \alpha^{1/3} x \end{aligned} \tag{1.2}$$

which transforms P_{II} to

$$w'' = 2w^3 + 1 + \frac{z}{\alpha} w, \tag{1.3}$$

where the primes denote z -derivatives. The three asymptotic regions are given by $z \ll \alpha$, $z \sim \alpha$, and $\alpha \ll z$. In this paper, we restrict ourselves to obtaining rigorous local asymptotic results in the first region. Note that this corresponds to $x \ll \alpha^{2/3}$.

These local results can be extended into the second region by using a nonlinear multiple-scales approach developed by Joshi and Kruskal [11, 5]

for the first and second Painlevé equations. The results give insight into a well known asymptotic limit of P_{II} . Consider [1] the following transformation of P_{II}

$$\begin{aligned} x &\mapsto \epsilon^2 X - \frac{6}{\epsilon^{10}} \\ y(x) &= \epsilon Y(X) + \frac{1}{\epsilon^5} \\ \alpha &= \frac{4}{\epsilon^{15}} \end{aligned}$$

In the limit as $\epsilon \rightarrow 0$, this leads to the first Painlevé equation (P_I)

$$Y'' = 6Y^2 + X. \quad (1.4)$$

It will be shown in a subsequent paper that this limit occurs at isolated points in the second region.

The method of Joshi and Kruskal was developed for the limit $x \rightarrow \infty$. The usual asymptotic studies of the Painlevé equations in the limit $x \rightarrow \infty$ apply here in the third region given by $\alpha^{2/3} \ll x$. To see this, consider the Boutroux transformation of P_{II} that makes the maximal dominant balance as $x \rightarrow \infty$ explicit, namely

$$\begin{aligned} y(x) &= \sqrt{x}u(t) \\ t &= \frac{2}{3}x^{3/2} \end{aligned} \quad (1.5)$$

under which P_{II} becomes

$$u_{tt} = 2u^3 + u + \frac{a - u_t}{t} + \frac{u}{9t^2}, \quad (1.6)$$

where $a = 2\alpha/3$. (See [5] for a deduction of Boutroux transformations based on maximal dominant balances.) The asymptotic analysis of this equation is usually carried out for $t \rightarrow \infty$. It remains valid so long as $a/t \ll 1$, i.e. $\alpha^{2/3} \ll x$.

1.2 Main Results

In section 2, we study the generic solutions of P_{II} . These are solutions defined by a large open set of initial data given at a point z_0 in the local region. Such

solutions are shown to have asymptotic behaviours given by elliptic functions as $\alpha \rightarrow \infty$. In section 3, we study limits of the generic solutions and their asymptotic behaviours. The union of the results of Sections 2 and 3 cover all solutions defined by all possible initial data at z_0 .

Moreover, the domain of validity of the generic locally valid elliptic-function-behaviour consists of the entire local domain $z \ll \alpha$ (punctured at poles and stationary points of the solution). Similarly, its degenerate limits are also valid in large subdomains of this region.

The solution in the nearly degenerate case is described by an almost-singly-periodic behaviour, exhibiting widely spaced lines of poles. There are further limits which exhibit only one line of poles or no such line within the domain of validity. We also consider the large energy limit and show that the resultant behaviours are still given by (scaled) elliptic functions in large subdomains.

Our results are given for both the solution and its first-derivative. By using P_{II} , these may be extended to yield the asymptotic behaviours of any higher-order derivative of $w(z)$.

In summary, we find all possible asymptotic behaviours of true solutions in the limit as $\alpha \rightarrow \infty$ in large domains lying in the region $z \ll \alpha$. We prove that these are asymptotic behaviours by finding explicit error bounds. We show that these behaviours are connected through limiting values taken by an energy-like parameter and are complete in the sense that they correspond to all possible values of initial data given at a point z_0 with $z_0 \ll \alpha$.

1.3 Notation

It should be noted that we use two symbols for an asymptotic balance between two functions $f(s)$ and $g(s)$ as $s \rightarrow \infty$ (along some path in the complex s -plane). First, if the limit of f/g is unity then we write $f \approx g$. Second, if a nonzero limit exists but differs possibly from unity, we write $f \sim g$. Our remaining usage of asymptotic notation is standard and definitions may be found, for example, in [24].

2 Generic Local Asymptotic Estimates

In this section, we prove that in the region where $z \ll \alpha$, the general solution of P_{II} is asymptotic to Jacobi elliptic functions. In particular, we estimate

the size of the asymptotic correction term. Moreover, we show that the spacing between zeroes of the general solution of P_{II} is closely approximated by a period of the locally valid elliptic function. It should be noted that we obtain an asymptotic result for both $w'(z)$ and $w(z)$. Note that we assume $|\alpha| > 1$ throughout.

First, we describe the local domain for our asymptotic analysis.

Definition 2.1 *Suppose $\epsilon \neq 0$, z_0 are given complex numbers and B is a positive real number. They are called appropriate if*

$$\begin{aligned} |\epsilon| &< 1, \frac{|\ln \epsilon|}{|\alpha|} < |\epsilon| \\ \left| \frac{z_0}{\alpha} \right| &< |\epsilon|, \frac{B}{|\alpha|} < 2|\epsilon| \\ (6 + 2\pi B)2\pi B &< |\epsilon|^{-5/7} |\ln \epsilon|^{-2} \end{aligned}$$

For an appropriate set $\{\epsilon, z_0, B\}$

$$\mathcal{Z} := \{z \mid |z - z_0| \leq B\},$$

is called the local z -domain.

Remarks:

1. The last inequality above is imposed to ensure that the error estimates obtained in Theorem 2.1 below is small. See further remarks following that theorem.
2. Note that one choice of appropriate $\{\epsilon, z_0, B\}$ is given by $\epsilon = 1/\sqrt{\alpha}$ with $z_0 \sim \alpha^{1/4}$, $B \sim \alpha^{1/8}$.
3. The above remark shows that z_0 and B do not have to be small. In particular, the local domain may be quite large (but not unboundedly large) even if z_0 is finite.

Standard theorems [25] show that for given bounded initial data $w(z_0)$, $w'(z_0)$ at a finite point z_0 there exists a unique locally analytic solution $w(z)$ of P_{II} . It is well known that $w(z)$ can be meromorphically extended [22] to any bounded domain. In particular, $w(z)$ is meromorphic in the local domain $\mathcal{Z}$.

We will work with an integrated form of P_{II} . Multiply Eqn(1.3) by w' and integrate as though the small term zw/α were not there. Then, we get

$$w'^2 = w^4 + 2w + 2E + 2 \int_{z_0}^z \frac{zww'}{\alpha} dz, \quad (2.7)$$

where E is a constant of integration. To leading-order, this equation is solved by Jacobian elliptic functions.

2.1 Elliptic Functions

To see that elliptic functions arise, and to fix the notation, let

$$P(w) := w^4 + 2w + 2E.$$

In general, $P(w)$ has four distinct roots, which we will denote by w_i , $i = 1, \dots, 4$, in the complex plane. The leading order equation

$$w'^2 = w^4 + 2w + 2E, \quad (2.8)$$

can be integrated again to give

$$\int_{\eta}^w \frac{dv}{\sqrt{P(v)}} = z - z_0, \quad (2.9)$$

where $\eta = w(z_0)$. The path of integration here lies on the Riemann surface defined by $\sqrt{P(v)}$. (This consists of two copies of the complex plane punctured at the roots of $P(v)$ and glued along lines connecting two pairs of roots.)

There exist two linearly independent closed contours C_j , $j = 1, 2$, on this surface each enclosing two roots of $P(v)$ in the v -plane. Paths connecting η to the same end-point w are deformable to each other modulo C_j . That is, deformations lead to Eqn (2.9) where w remains unchanged while the right side changes by integer multiples of

$$\omega_j = \oint_{C_j} \frac{dv}{\sqrt{P(v)}}. \quad (2.10)$$

Further properties of the elliptic integrals ω_j , including differential equations they satisfy as functions of E , are given in Appendix A.

This shows that the solution $w(z)$ (the inverse of the function $z(w)$ defined by Eqn(2.9)) has two periods given by ω_j . Moreover, it is straightforward to show that $w(z)$ is meromorphic. That is, it is an elliptic function.

One of the two periods of w becomes infinite (while the other remains finite) at special values of E . These are the values for which two roots of $P(w)$ coincide. To find these, consider

$$P'(w) = 4w^3 + 2 = 0 \Rightarrow w = d_k := (-1/2)^{1/3}, k = 1, 2, 3. \quad (2.11)$$

If such points are also zeroes of $P(w)$ then we must have

$$E = D_k := -\frac{3}{4}(-1/2)^{1/3} = \left(\frac{27}{128}\right)^{1/3}, k = 1, 2, 3. \quad (2.12)$$

For each such value of E , $P(w)$ has a double root. Note, however, that $P(w)$ cannot have a triple root, because $P''(w) = 12w^2$ cannot vanish simultaneously with $P'(w)$.

Suppose $E = D_k$, for some $k = 1, 2$, or 3 . Let ρ and σ be the roots of $P(w)$ distinct from d_k . Then rewriting

$$P(w) = (w - d_k)^2(w - \rho)(w - \sigma)$$

in expanded form shows that

$$\rho\sigma = -\frac{3}{2d_k}, \quad \rho + \sigma = \frac{1}{d_k^2}.$$

Solving these equations, we get

$$\rho, \sigma = \frac{1}{2d_k^2} \{1 \pm i\sqrt{2}\}.$$

Note that these cannot equal d_k because $P(w)$ cannot have a triple root. That is, there exists $c > 0$, such that $|\rho - d_k| > c$, $|\sigma - d_k| > c$.

Now suppose $E = D_k + \delta^2$, for $\delta \ll 1$. Then setting $P(w) = 0$ and using the fact that $d_k - \rho$, $d_k - \sigma$ are bounded below, it is straightforward (see [26] for detailed proofs of similar results) to show that the two nearly coalescing roots w_j of $P(w)$ satisfy

$$w_j = d_k + O(\delta),$$

and that the other two roots are a finite distance away from d_k .

Note also that if E is large, say $|E| = \log \delta$, then the periods become small. To see this, consider the roots of $P(w)$. The roots now have modulus at least equal to $O(|E|^{1/4})$ and so

$$\omega_j = O(|E|^{-3/4}), j = 1, 2.$$

(Such asymptotic results for the complete elliptic integrals can be obtained straightforwardly from the differential equations derived in Appendix A.) Hence the poles of the elliptic function crowd together and the function itself becomes large even at points of holomorphy.

2.2 Generic Solutions

Our main result holds for solutions that are described by a large open and dense set of initial values that we call *generic*.

Definition 2.2 *Suppose ϵ is an appropriate given number. Parameters η, η' are called generic if for*

$$2E = \eta'^2 - \eta^4 - 2\eta,$$

they satisfy

$$\begin{aligned} |\epsilon|^{1/7} &< |\eta'| < |\ln \epsilon|^2, \quad |\eta| < |\ln \epsilon| \\ 2|\epsilon|^{2/7} &< |E - D_k| \end{aligned}$$

These supply the initial values for the solutions and ensure that their formal elliptic-function-type behaviours are not too close to degenerate ones. Initial values that violate this definition are studied in section 3.

Definition 2.3 *Suppose ϵ, z_0 are appropriate numbers. Let $w(z)$ be a solution of P_{II} defined by generic initial values η, η' at z_0 , i.e. $w(z_0) = \eta, w'(z_0) = \eta'$. Then $w(z)$ is called a generic solution.*

To get valid asymptotic results, we need to confine such a solution $w(z)$ to domains that avoid its poles and stationary points.

Definition 2.4 *Given an appropriate ϵ , define the w -domain by*

$$\mathcal{W} := \{w \mid |w| < |\ln \epsilon|, |P(w)| > |\epsilon|^{1/7}\}$$

and z -domain by

$$\tilde{\mathcal{Z}} := \{z \in \mathcal{Z} \mid w(z) \in \mathcal{W}\}.$$

ϵ is called small enough if $\tilde{\mathcal{Z}}$ is connected.

The direct proof [22] of the Painlevé property of P_{II} shows that there exists an open disk of values ϵ for which $\tilde{\mathcal{Z}}$ is connected. (Explicit lower bounds for $A := 1/|\epsilon|$, may be found on p. 197 of [22].) Moreover, any two points in $\tilde{\mathcal{Z}}$ are connected by paths γ , of bounded length $2\pi B$, such that their images Γ under w lie in $\mathcal{W}$ with length $< 2\pi |\ln \epsilon|$.

Our major result is

Theorem 2.1 *Given an appropriate set $\{\epsilon, z_0, B\}$ where ϵ is small enough, and generic initial values η, η' , the generic solution $w(z)$ satisfies*

$$\begin{aligned} |w'^2 - P(w)| &\leq \frac{\kappa}{2} |\epsilon| |\ln \epsilon|^2 \\ \left| \int_{\eta}^w \frac{dv}{\sqrt{P(v)}} - (z - z_0) \right| &\leq 2\pi \kappa B |\epsilon|^{6/7} |\ln \epsilon|^2 \end{aligned}$$

for all $z \in \tilde{\mathcal{Z}}$, where $\kappa := 6 + 2\pi B$. Moreover, if z_0 and z_{10} are two successive points in $\tilde{\mathcal{Z}}$ where $w = \eta$, then for $j = 1$ or 2 ,

$$|(z_{10} - z_0) - \omega_j| \leq 2\pi \kappa B |\epsilon|^{6/7} |\ln \epsilon|^2.$$

Proof: Equation(2.7) can be written

$$(w'(z))^2 = P(w(z)) + 2S$$

where

$$S := \int_{\gamma} \frac{z}{\alpha} w w' dz \quad (2.13)$$

where γ is a path connecting z_0 to z in $\tilde{\mathcal{Z}}$. The assumptions give

$$\left| \frac{z}{\alpha} \right| \leq \frac{B}{|\alpha|} + \left| \frac{z_0}{\alpha} \right| < 3|\epsilon|.$$

Integration by parts shows that

$$S = \left[\frac{zw^2}{2\alpha} \right]_{(z_0, \eta)}^{(z, w)} - \frac{1}{2\alpha} \int w^2 dz.$$

Hence we get

$$\begin{aligned} 2|S| &\leq 6|\epsilon| |\ln \epsilon|^2 + |\epsilon| \cdot 2\pi B |\ln \epsilon|^2 \\ &= \kappa |\epsilon| |\ln \epsilon|^2 \end{aligned}$$

as desired. Now integrate Eqn(2.7) once more by taking the square root and dividing by $\sqrt{P(w)}$. We get

$$\int_{\eta}^w \frac{dv}{\sqrt{P(v)}} = \int_{\gamma} dz \left\{ 1 + \frac{2S}{P(w)} \right\}^{1/2}. \quad (2.14)$$

Note that we fix a choice of branch of the square root here, so that as $\epsilon \rightarrow 0$ we have the positive real branch. From Eqn(2.14) we get

$$\int_{\eta}^u \frac{dv}{\sqrt{P(v)}} - (z - z_0) = \int_{z_0}^z dz \left\{ \left(1 + \frac{2S}{P(w)} \right)^{1/2} - 1 \right\}. \quad (2.15)$$

Note that

$$\left| \frac{2S}{P(w)} \right| \leq \kappa |\epsilon|^{6/7} |\ln \epsilon|^2. \quad (2.16)$$

Let

$$Q := \left(1 + \frac{2S}{P(w)} \right)^{1/2} + 1.$$

Then we have

$$Q^2 = 2Q + \frac{2S}{P(w)} \Rightarrow Q = 2 + \frac{2S}{QP(w)}.$$

The choice of branch ensures that Q is lower bounded by $|Q| \geq 1$. Therefore, it follows that

$$|Q - 2| \leq \left| \frac{2S}{P(u)} \right| \Rightarrow \left| \sqrt{1 + \frac{2S}{P(u)}} - 1 \right| \leq \kappa |\epsilon|^{6/7} |\ln \epsilon|^2.$$

Integration then gives the desired result. The separation $z_{10} - z_0$ is based on the same result with the path of integration Γ being a closed contour enclosing two roots of $P(u)$. $\square$

Remarks:

1. The crucial element of this proof lies in the lower bound for $P(w)$. The microscopic details of this lower bound can be changed without affecting the macroscopic form of the asymptotic result. For example, suppose the conditions on η , η' are changed to allow $E - D_k = \delta^2$ where $|\delta| \sim |\epsilon|^{p/7}$,

$p < 7$. Then the same result as in the theorem holds for w'^2 , and the second (and third) results of the theorem hold with the order of the error estimate changed to

$$|\epsilon|^{1-p/7} |\ln \epsilon|^2.$$

2. It should be noted, however, that if δ is small enough (but nonzero), then the incomplete elliptic integral on the left of the first result is more accurately written as the inverse of a hyperbolic function. See the next section for details.
3. The third result of the theorem holds for any two successive points (in a direction given by ω_j for a chosen j) at which w takes the same value. By taking ϵ to be smaller, such points can be taken arbitrarily close to poles of w . Hence the result also holds for the spacing between two poles.
4. Notice that the incomplete elliptic integral

$$\int_{\eta}^w \frac{dv}{\sqrt{P(v)}}$$

is still convergent if the generic initial value η happens to be arbitrarily close to a zero of $P(w)$. However, if it is so close to such a zero that the path intrudes into a “hole” of $\tilde{\mathcal{Z}}$, then we need a slightly modified argument for the proof. In this case, evaluating P_{II} at z_0 gives a finite nonzero lower-bound for $|w''(z_0)|$. Therefore, there exists an ascent path for $|w'|$ starting at z_0 . Following this path (over a bounded distance) takes the solution to a point z_1 where $P(w)$ is lower-bounded by $|\epsilon|^{1/7}$. We can now start again at such a point and apply the theorem. Such arguments are used in more detail in section 3.

5. In the proof above, B is not required to be finite, but it should be *appropriate*. As remarked after Definition 2.1, B may be quite large. For example, the error estimates of the theorem are still small for $B \approx \epsilon^{-1/4} \approx \alpha^{1/8}$.

The last remark above shows that for any finite z_0 and large but appropriate B , a generic solution w has the *same* elliptic-function-behaviour throughout the “local” domain $\tilde{\mathcal{Z}}$. The error, however, does increase with B . If B increases beyond appropriate values, z encroaches on the second asymptotic region (i.e. $z \sim \alpha$) which will form the subject of a subsequent study.

3 Limits of Generic Solutions

In this section, we consider the possible asymptotic behaviours of generic solutions $w(z)$ when the parameter E or initial values η, η' become close to degenerate values or become large.

Degenerate limits occur when $E - D_k =: \delta^2$, with $|\delta| \ll 1$, for some $k = 1, 2$ or 3 . We separate this into the following subcases.

1. For some integer $p > 1$, $|\epsilon|^{p/7} < |\delta| \leq |\epsilon|^{1/7}$
2. $|\delta| \leq |\epsilon|^{p/7}$, for any integer $p > 1$
3. For some $k = 1, 2$ or 3 , $|\eta'| \leq |\epsilon|^{1/7}$, $|\eta - d_k| \leq |\epsilon|^{1/14}$

Note that these cases overlap. The first subcase is studied in Subsection 3.1 and the remaining subcases are studied in Subsection 3.2 below.

The results of the direct proof of the Painlevé property of P_{II} [22] show that large initial values with finite E imply that the initial point z_0 is close to a pole of the generic solution $w(z)$. The proof is a trivial rewriting of the proof given in section 3 of [22] and so we do not reproduce it here. In this case, Theorem 2.1 shows that the pole of the Painlevé transcendent nearest to z_0 lies close to a pole of the elliptic-function-type asymptotic behaviour of $w(z)$.

Therefore, the limit of interest, that violates the genericity condition, is when E becomes unbounded. (Equivalently, we may take η, η' large with $\eta' \neq \eta^4$.) In Subsection 3.3, we consider the large energy limit when $|E| \geq |\ln \epsilon|^4$.

3.1 Nearly Degenerate Solution

The results of Theorem 2.1 continue to apply even if δ is smaller than $\epsilon^{1/7}$. (See Remark 1 following Theorem 2.1.) However, for small enough δ , the elliptic-function-type behaviours of $w(z)$ are more accurately represented by singly periodic behaviours. We show this here.

From the results of Section 2.1, we have that if $E = D_k + \delta^2$, where $\delta \ll 1$,

$$P(w) = (w - d_k)^2(w - \rho)(w - \sigma) + 2\delta^2.$$

Recall $d_k = (-1/2)^{1/3}$, and ρ, σ equals one of

$$\frac{1}{2d_k^2} \{1 \pm i\sqrt{2}\}.$$

Definition 3.1 Let $\{\epsilon, z_0, B\}$ be an appropriate set, $k = 1, 2$, or 3 , and $1 < p$ be a given number. Parameters η, η' are called nearly degenerate if for

$$2E = \eta'^2 - \eta^4 - 2\eta, \quad E - D_k =: \delta^2,$$

they satisfy

$$|\epsilon|^{1/7} < |\eta'| < |\ln \epsilon|^2, \quad |\epsilon|^{1/14} < |\eta - d_k| < |\ln \epsilon|$$

$$|\epsilon|^{p/7} < |\delta| \leq |\epsilon|^{1/7}$$

Note that the condition on E implies η', η lie in a one-complex dimensional deleted neighbourhood which excludes the point $\eta' = 0, \eta = d_k$, for a given k . This deleted neighbourhood of initial conditions gives rise to a solution that is closer to a singly periodic limit of the previous elliptic function.

Definition 3.2 Suppose ϵ, z_0 are appropriate numbers and η, η' are nearly degenerate parameters. Then the solution $w(z)$ of P_{II} given by $w(z_0) = \eta, w'(z_0) = \eta'$ is called a nearly degenerate solution.

Now the appropriate polynomial to consider is

$$R(w) := (w - d_k)^2(w - \rho)(w - \sigma),$$

and the $\mathcal{W}$ -domain becomes

$$\mathcal{V} = \{w \mid |w| < |\ln \epsilon|, |R(w)| > \epsilon^{1/7}\}.$$

Correspondingly, ϵ is called small enough if the preimage $\tilde{\mathcal{Z}}_\delta$ of $\mathcal{V}$ in $\mathcal{Z}$ is connected. To be specific, we take $k = 1$, with real

$$d_1 = -1/2^{1/3}.$$

The main result in this case is

Theorem 3.1 Given an appropriate set $\{\epsilon, z_0, B\}$, with $\kappa|\epsilon|^{5/7}|\ln \epsilon|^2 \leq 1$, where ϵ is small enough, and nearly degenerate parameters η, η' , the nearly degenerate solution satisfies

$$|w'^2 - R(w)| < 3|\epsilon|^{2/7},$$

$$\left| \int_\eta^w \frac{dv}{(v - d_1)\sqrt{(v - \rho)(v - \sigma)}} - (z - z_0) \right| < 6\pi B|\epsilon|^{1/7},$$

for all $z \in \tilde{\mathcal{Z}}_\delta$.

Proof: The proof is almost identical to that of Theorem 2.1 with a modified small term. Note that

$$w'^2 = R(w) + 2\delta^2 + 2S,$$

where S is as defined in Eqn (2.13). Therefore, since we still have

$$2|S| \leq \kappa|\epsilon| |\ln \epsilon|^2$$

whereas

$$|\delta|^2 \leq |\epsilon|^{2/7},$$

we get the first result. Integrating once more, we get

$$\int_{\eta}^w \frac{dv}{\sqrt{R(v)}} = \int_{z_0}^z dz \left\{ 1 + \frac{2\delta^2}{R(w)} + \frac{2S}{R(w)} \right\}^{1/2}. \quad (3.17)$$

By assumption, we have

$$\left| \frac{2\delta^2}{R(w)} \right| < 2|\epsilon|^{1/7}$$

and

$$\left| \frac{2S}{R(w)} \right| < \kappa|\epsilon|^{6/7} |\ln \epsilon|^2.$$

So we have

$$\left| \frac{2\delta^2}{R(w)} + \frac{2S}{R(w)} \right| < 3|\epsilon|^{1/7}.$$

The remainder of the proof of Theorem 2.1 can be repeated with $\{2\delta^2 + 2S\}/R(w)$ in place of $2S/P(w)$ and $R(w)$ in place of $P(w)$ to get the desired result. $\square$

Note that we now require a smaller B to get asymptotic validity. Namely, we need $6\pi B\epsilon^{1/7} < 1$. For example, $6\pi B < |\epsilon|^{-1/14}$ would be sufficient.

The interesting difference from the generic case arises because the integral on the left can be evaluated explicitly in terms of hyperbolic functions. To see this, make the change of variables:

$$v = \frac{(\sigma - \rho)}{2} \left(\frac{f + f^{-1}}{2} \right) + \frac{(\sigma + \rho)}{2} \quad (3.18)$$

$$f = \sqrt{3}g - \sqrt{2}i \quad (3.19)$$

Then the integral becomes

$$\int_{\eta}^w \frac{dv}{(v-d_1)\sqrt{(v-\rho)(v-\sigma)}} = \frac{2}{d_1\sqrt{6}}i \int \frac{dg}{g^2+1}.$$

Hence we get

$$g = i \tanh \left\{ \frac{\sqrt{6}d_1}{2}(z-z_0) + a_0 + O(\epsilon^{1/7}) \right\}, \quad (3.20)$$

where $a_0 = \tanh^{-1}(g(z_0))$. Note that this behaviour is singly periodic and possess a line (rather than a lattice) of poles. In a direction orthogonal to the line, the behaviour becomes asymptotic to constants.

There is an alternative change of variables. Consider the transformation:

$$v = d_1 + u \quad (3.21)$$

$$u = -1/F \quad (3.22)$$

$$F = \frac{1}{\sqrt{18d_1^2}}G + \frac{1}{3d_1} \quad (3.23)$$

Then the integral becomes

$$\int_{\eta-d_1}^{w-d_1} \frac{du}{u\sqrt{u^2+4d_1u+6d_1^2}} = \frac{1}{\sqrt{6}d_1} \int \frac{dG}{\sqrt{G^2+1}}.$$

In this case, we get

$$G = \sinh \left\{ \sqrt{6}d_1(z-z_0) + b_0 + O(\epsilon^{1/7}) \right\}, \quad (3.24)$$

where $\sinh b_0 = G(z_0) = -\sqrt{2} + \sqrt{18}d_1/(d_1 - \eta)$. The behaviours indicated by Eqns(3.20) and (3.24) are clearly related and represent the same nearly degenerate solution in the domain $\tilde{\mathcal{Z}}_{\delta}$. Note that one line of poles is given approximately by

$$z_0 + \frac{-b_0 + 2\pi ni}{\sqrt{6}d_1} \quad (3.25)$$

for integer n .

In Appendix A, we show that one of the periods ω_j of the generic elliptic-function behaviour becomes logarithmically large and one remains finite as

$E \rightarrow D_k$ for some $k = 1, 2, 3$. For the nearly degenerate case this shows that for some j

$$\omega_j \sim \ln |E - D_k| \sim |\ln \epsilon|.$$

For sufficiently small ϵ , this is clearly smaller than B (e.g. $B \sim |\epsilon|^{-1/14}$) for which the asymptotic results of Theorem 3.1 are valid. Therefore, the results above suggest that more than one line of poles may be visible in the domain $\tilde{\mathcal{Z}}_\delta$. Between the lines of poles, the local behaviour is still described by Eqn (3.24).

3.2 Degenerate Solution

Here we consider the case when δ is arbitrarily small or when initial values are arbitrarily close to degenerate values.

Definition 3.3 *Let $\{\epsilon, z_0, B\}$ be an appropriate set, and, for some $k = 1, 2, 3$, let η, η' be given parameters.*

- *If they satisfy*

$$2E := \eta'^2 - \eta^4 - 2\eta = 2D_k + 2\delta^2,$$

where $|\delta| \leq |\epsilon|^{p/7}$ for all integers $p > 1$, and

$$|\epsilon|^{1/7} < |\eta'| < |\ln \epsilon|^2, \quad |\epsilon|^{1/14} < |\eta - d_k| < |\ln \epsilon|$$

then they are called almost degenerate parameters.

- *If, on the other hand, they satisfy*

$$|\eta'| \leq |\epsilon|^{1/7}, \quad |\eta - d_k| \leq |\epsilon|^{1/14}$$

then they are called degenerate parameters.

To be concrete, we take $k = 1$. Assume that $\epsilon, \mathcal{V}, \tilde{\mathcal{Z}}_\delta$ are as defined in the previous section, with a slight modification that $|R| \geq |\epsilon|^{1/7}/16$ on $\tilde{\mathcal{Z}}_\delta$.

Definition 3.4 *Let ϵ be small enough such that the domain $\tilde{\mathcal{Z}}_\delta$ is connected and let η, η' be almost degenerate parameters. Then the solution $w(z)$ of P_{II} given by $w(z_0) = \eta, w'(z_0) = \eta'$ is called an almost degenerate solution.*

The solution defined by almost degenerate parameters has asymptotic behaviour that is given locally by a singly periodic function. In fact, the proof of the last subsection can be applied to show that the following theorem holds.

Theorem 3.2 *Given an appropriate set $\{\epsilon, z_0, B\}$, with $\kappa|\epsilon|^{5/7}|\ln \epsilon|^2 \leq 1$, where ϵ is small enough, and almost degenerate parameters η, η' , the almost degenerate solution satisfies*

$$|w'^2 - R(w)| < \frac{\kappa}{2}|\epsilon| |\ln \epsilon|^2$$

$$\left| \int_{\eta}^w \frac{dv}{(v - d_1)\sqrt{(v - \rho)(v - \sigma)}} - (z - z_0) \right| < 2\pi\kappa B|\epsilon|^{6/7}|\ln \epsilon|^2.$$

for all $z \in \tilde{\mathcal{Z}}_{\delta}$.

Note that the domain of validity is now larger than in Theorem 3.1, i.e. B may be as large as $1/|\epsilon|^{5/7}$.

The almost degenerate case clearly also has representations given by Eqns(3.20) and (3.24) in the domain $\tilde{\mathcal{Z}}_{\delta}$. However, now there may be at most one line of poles. The reason lies in the fact that the almost degenerate limit of an elliptic-function behaviour has a period that is possibly $\gg |\ln \epsilon|$. For example, suppose $\delta \sim \exp(-1/|\epsilon|)$. Then

$$\omega_j \sim \ln(E - D_k) \sim 1/|\epsilon|.$$

Hence the distance between two lines of poles (or any two lines formed by interpolation through integer multiples of the finite period) may be much larger than appropriate values of B .

Nevertheless, there must be at least one line of poles in the local domain, because the lower bound on $\eta - d_1$ implies $|b_0| < |\ln \epsilon|$. Hence the sequence given by Eqn(3.25) lies in $\tilde{\mathcal{Z}}_{\delta}$.

The remaining case, apparently different to this, is the degenerate case. However, we can show that the almost degenerate solution becomes a degenerate solution somewhere in its domain of validity. To show this, we travel on a descent path. Recall that a complex-valued function $f(z)$ of one complex variable z possesses a descent path (for its modulus) starting a point z_0 if $f'(z_0) \neq 0$. Along the steepest descent path, we have $|df| = -d|f|$.

Theorem 3.3 *Let $\{\epsilon, z_0, B\}$ be an appropriate set with $|\epsilon|^{1/4} < 2^{-1/3}$. Let η, η' be almost degenerate parameters and let w be an almost degenerate solution. Then there exists a path Γ joining z_0 to a point z_1 where $|w - d_k|^2$ and $|w'|$ become less than $|\epsilon|^{1/7}$, for some $k = 1, 2, 3$.*

Proof: *Case 1:* Consider the case when $|\eta| \geq |d_k|/4$. By assumption of appropriateness and upperbounds on η, η' , we have

$$\left| \frac{\eta + z_0 \eta'}{\alpha} \right| < 2|\epsilon| |\ln \epsilon|^2.$$

Therefore, we can lower-bound

$$w''' = 6w^2 w' + \frac{1}{\alpha}(w + zw') \quad (3.26)$$

at (and by continuity near) z_0 by

$$|w'''(z_0)| > |d_k|^2 |\epsilon|^{1/7} / 4.$$

In other words, there exists a descent path for w'' starting at z_0 . We take Γ to be this descent path.

Recall that in P_{II}

$$w'' = 2w^3 + 1 + \frac{z}{\alpha}w \quad (3.27)$$

and its first integral

$$w'^2 = R(w) + 2S, \quad (3.28)$$

we have

$$\left| \frac{z}{\alpha}w \right| < 3|\epsilon| |\ln \epsilon|, \quad |S| < \kappa/2 |\epsilon| |\ln \epsilon|^2 \quad (3.29)$$

throughout $\tilde{\mathcal{Z}}_\delta$, and in particular on the part of the path Γ lying in $\tilde{\mathcal{Z}}_\delta$.

Since w'' is decreasing, this shows that $2w^3 + 1$ must decrease along Γ , until it reaches a first point z_1 where $|w - d_k|$ becomes smaller than $|\epsilon|^{1/14}$, or where we reach the boundary of $\tilde{\mathcal{Z}}_\delta$. Similarly $R(w)$ also decreases along Γ .

At that point, the definitions of d_k, ρ , and σ show that

$$|R(w(z_1))| \geq \frac{|\epsilon|^{1/7}}{16},$$

while it remains larger than this bound earlier on Γ . Moreover, we can estimate its length. For conciseness, we rename $v := w''$. By definition we get

$$\begin{aligned} z_1 - z_0 &= \int_{z_0}^{z_1} dz \\ &= \int_{\Gamma} \frac{dv}{v'} \end{aligned}$$

But from Eqns (3.26), (3.28), and (3.29), we get

$$|v'| \geq 4|w|^2|w'| \geq |d_k|^2|\epsilon|^{1/14}/64$$

if

$$|\epsilon|^{1/14} > 32\pi^2 d^2 |\epsilon| |\ln \epsilon| \quad (3.30)$$

where d is the length of Γ . We then get

$$\begin{aligned} d &= \left| \int_{\Gamma} \frac{dv}{v'} \right| \\ &\leq \int_{\Gamma} \frac{|dv|}{|v'|} \\ &\leq -\frac{64}{|d_k|^2} \int_{\Gamma} d|v|/|\epsilon|^{1/14} \\ &= \frac{64|\epsilon|^{-1/14}}{|d_k|^2} (|v(z_0)| - |v(z_1)|) \end{aligned}$$

This clearly satisfies Eqn(3.30). Moreover, this shows that Γ lies within $\tilde{Z}_{\delta}$.

Case 2: Now consider the case when $|\eta| < |d_k/4|$. We have from

$$w'' = 2w^3 + 1 + \frac{z}{\alpha}w$$

that

$$|w''(z_0)| \geq 1/2.$$

Hence there exists a descent path for w' starting at z_0 . We take Γ to be this path. The decrease in $|w'|$ implies a corresponding decrease in $|R(w)|$. Therefore, w must get closer to a root of $R(w)$. Since all such roots have modulus $\geq 1/2^{1/3}$, $|w|$ must increase. There exists a first point $\tilde{z}_0$ on Γ where w has modulus equal to $|d_k/4|$. That is,

$$|w(\tilde{z}_0) - d_k| \geq \frac{3|d_k|}{4}.$$

At such a point, we have

$$|R(w)| > 18|d_k|^2,$$

by using the definitions of d_k , ρ , and σ . Using arguments similar to above we get

$$|\tilde{z}_0 - z_0| =: d \leq \frac{1}{2\sqrt{18}}.$$

We now use the results of Case 1 starting at $\tilde{z}_0$ to extend Γ to z_1 . $\square$

It is instructive to consider the limits of Theorems 3.1, 3.2, explicitly when

$$0 \neq |w - d_k| < |\epsilon|^{1/14}, \quad 0 \neq |\eta - d_k| < |\epsilon|^{1/14}.$$

We have

$$\begin{aligned} & \int_{\eta}^w \frac{dv}{\sqrt{R(w)}} \\ &= \frac{1}{\sqrt{6}d_k} \int_{\eta}^w \frac{dv}{(w - d_k) \sqrt{1 + 2(w - d_k)/(3d_k) + (w - d_k)^2/(6d_k^2)}} \\ &= \frac{1}{\sqrt{6}d_k} \int_{\eta}^w \frac{dv}{(w - d_k)} \left\{ 1 - (w - d_k)/(3d_k) + O((w - d_k)^2) \right\} \\ &= \frac{1}{\sqrt{6}d_k} \left\{ \ln((w - d_k)/(\eta - d_k)) - (w - \eta)/(3d_k) + O((w - d_k)^2) \right\} \end{aligned}$$

Taking the inverse of the second result of Theorem 3.2 implies that

$$\begin{aligned} w - d_k &= (\eta - d_k) \exp\left(\sqrt{6}d_k(z - z_0)\right) \left(1 - (\eta - d_k)/(3d_k) + O(|\epsilon|^{1/7})\right) \\ &+ \frac{1}{3d_k}(\eta - d_k)^2 \exp\left(2\sqrt{6}d_k(z - z_0)\right) (1 + O(|\epsilon|^{1/14})). \end{aligned} \quad (3.31)$$

It is clear that the exponential terms here will grow in a half-plane of directions in $\tilde{\mathcal{Z}}_{\delta}$. For example, consider $d_1 = -1/2^{1/3}$. Then

$$\Re(z - z_0) < 0 \Rightarrow \Re(d_1(z - z_0)) > 0.$$

In this half-plane, the solution grows away from the initial assumed value. By following an ascent path, it is straightforward to show that this behaviour matches with that given by Eqn(3.24) in this half-plane.

Conversely, if the solution has degenerate initial values, we can integrate P_{II} to show that w is given by Eqn(3.31) in a local domain around z_0 . To prove this, we consider a domain given by

$$\mathcal{U} = \{w \mid |\epsilon|^{1/7} < |w| < |\ln \epsilon|, |w - d_k| < \epsilon^{1/14}\}.$$

Correspondingly, define ϵ to be small enough if the preimage $\tilde{\mathcal{Z}}_0$ of $\mathcal{U}$ in $\mathcal{Z}$ is connected.

Definition 3.5 Let ϵ be small enough and let η, η' be degenerate parameters. Then the solution $w(z)$ of P_{II} given by $w(z_0) = \eta, w'(z_0) = \eta'$ is called a degenerate solution.

Then we have the following theorem.

Theorem 3.4 Assume we are given an appropriate set $\{\epsilon, z_0, B\}$, where ϵ is small enough such that

$$\kappa|\epsilon|^{5/7}|\ln \epsilon|^2 \leq 1, \quad |\epsilon|^{1/7} < |d_k|,$$

and

$$6|d_k|^2 + 4|d_k||\epsilon|^{1/14} + |\epsilon|^{1/7} < 7|d_k^2|.$$

Furthermore, assume that degenerate parameters $\eta \neq d_k, \eta' \neq 0$, are given. Then there exists a number $\mu > 0$ such that the degenerate solution satisfies

$$|w'^2 - 6d_k^2(w - d_k)^2| \leq 14|d_k|^2|\epsilon|^{1/7},$$

$$\left| \int_{\eta}^w \frac{dv}{\sqrt{6d_k^2(v - d_k)^2 + 2\delta^2}} - (z - z_0) \right| < 2\pi B\mu|\epsilon|^{1/14}.$$

for all $z \in \tilde{\mathcal{Z}}_0$.

Proof: Again, the technique of the proof is almost the same as Theorem 2.1 and we only highlight the important differences. After one integration, we get

$$w'^2 = R(w) + 2\delta^2 + 2S,$$

where by the given assumptions, we have

$$2|\delta|^2 = 2|E - D_k| \leq 8|d_k|^2|\epsilon|^{1/7}.$$

S here is as before, with the same upperbound: $|2S| < \kappa|\epsilon||\ln \epsilon|^2$. To get the first result, we use the fact that

$$R(w) = 6d_k^2(w - d_k)^2 + 4d_k(w - d_k)^3 + (w - d_k)^4,$$

and the bounds on $w - d_k$.

Hence it is slightly more transparent to write

$$w'^2 = 6d_k^2(w - d_k)^2 + 4d_k(w - d_k)^3 + (w - d_k)^4 + 2\delta^2 + 2S.$$

The second result follows from taking the square root, dividing by $\sqrt{6d_k^2(w - d_k)^2 + 2\delta^2}$ and bounding the remaining error terms. $\square$

Remarks:

1. Note that in the case $\eta = d_k$, $\eta' = 0$, $E = D_k$ and the integral in the second result above is divergent. We consider this case here. At z_0 , we have

$$w''(z_0) = z_0\eta/\alpha, w'''(z_0) = \eta/\alpha,$$

and so there exists an ascent path for w'' starting at z_0 . By P_{II}, this is also an ascent path for $2w^3 + 1$. Following this path, even for a short distance (say of order $\epsilon^{1/14}$) allows us to reach a point z_1 , say, where we can use the above argument for small but nonzero $w(z_1) - d_k$.

2. The above theorem extends easily to the case when $\eta \neq d_k$ but $\eta' = 0$, because $E - D_k \neq 0$.
3. The case when $\eta = d_k$, but $\eta' \neq 0$ can be treated by an argument similar to the above ascent-path argument.
4. The only remaining difficulty is that the path of integration may originate at or possibly intersect one of the simple zeroes of $R(w)$. (Note that the definition of $\mathcal{U}$ does not exclude the simple zeroes ρ, σ of $R(w)$ from the path.) However, these problems are clearly overcome by ascent-path-type argument similarly to the way in which the zeroes of $P(w)$ were avoided in Remark 4 following the proof of Theorem 2.1.

These remarks cover all the remaining degenerate possibilities.

3.3 Large Energy Limit

Here we consider the only remaining case of possible limits of the generic solution, that is, we consider the case when $|E|$ (or equivalently η or η' , with $\eta' \neq \eta^4$) is large.

Definition 3.6 *Suppose ϵ is an appropriate number. Parameters η, η' are called large if*

$$2E = \eta'^2 - \eta^4 - 2\eta$$

satisfies

$$|E| \geq |\ln \epsilon|^4.$$

The corresponding solution w defined by initial values $w(z_0) = \eta$ $w'(z_0) = \eta'$ is called a large solution.

In this case, it is convenient to rescale variables as follows. Let

$$\begin{aligned} w &:= E^{1/4} q(s) \\ s &:= E^{1/4} z \end{aligned}$$

Then the first integral of P_{Π} becomes

$$q_s^2 = q^4 + 2 + \frac{2}{E^{3/4}} \left(q + \int ds s q q_s / \alpha \right). \quad (3.32)$$

Definition 3.7 *Define the q -domain*

$$\mathcal{Q} := \left\{ sE^{-1/4} \in \mathcal{Z} \mid |q| < |E|^{1/4}, |q^4 + 2| > |E|^{-1/4} \right\}$$

and s -domain

$$\tilde{\mathcal{S}} := \{ sE^{-1/4} \in \mathcal{Z} \mid q(s) \in \mathcal{Q} \}.$$

ϵ is called small enough if $\tilde{\mathcal{S}}$ is connected.

Let $s_0 = z_0 E^{1/4}$, $\iota = \eta E^{-1/4}$, $\iota' = \eta' E^{-1/2}$. We will loosely call ι , ι' , and $q(s) = E^{-1/4} w(sE^{-1/4})$ large if η , η' and $w(z)$ are large.

Note that we will now restrict paths of integration to those with length upperbounded by $2\pi B$ in $\tilde{\mathcal{S}}$.

Theorem 3.5 *Given an appropriate set $\{\epsilon, z_0, B\}$, with $|\epsilon\kappa|/2 < 1/2$ where ϵ is small enough, and large ι , ι' , and the corresponding solution $q(s)$ satisfies*

$$\begin{aligned} \left| q_s^2 - (q^4 + 2) \right| &\leq \frac{3}{2} |E|^{-1/2} \\ \left| \int_{\iota}^q \frac{dq}{\sqrt{q^4 + 2}} - (s - s_0) \right| &\leq 3\pi B |E|^{-1/2} \end{aligned}$$

for all $s \in \tilde{\mathcal{S}}$. Moreover, if s_0 and s_{10} are two successive points in $\tilde{\mathcal{S}}$ where $w = \iota$, then for $j = 1$ or 2 ,

$$|(s_{10} - s_0) - \omega_j| \leq 3\pi B |E|^{-1/2}.$$

Proof: The small term in the first integral Eqn(3.32) is given by

$$T := \frac{1}{E^{3/4}} \left(q + s q^2 / (2\alpha) - \int ds q^2 / (2\alpha) \right).$$

The results follow from the techniques of the proof of Theorem 2.1. $\square$

4 Acknowledgements

It is a pleasure to thank the Isaac Newton Institute, where this study was first started, and the Australian Research Council for their support.

A Complete Elliptic Integrals

Here we prove some results for complete elliptic integrals as functions of E .

Let C be one of C_k , $k = 1, 2$.

Lemma A.1 *The elliptic integrals*

$$\tilde{\psi} = \oint_C dw w^2 \sqrt{w^4 + 2w + 2E},$$

$$\psi = \oint_C dw \frac{w^2}{\sqrt{w^4 + 2w + 2E}},$$

$$\psi' = - \oint_C dw \frac{w^2}{w^4 + 2w + 2E^{3/2}},$$

considered as functions of E satisfy the following differential equations.

$$\begin{aligned} \frac{d\tilde{\psi}}{dE} &= \psi \\ \frac{d\psi}{dE} &= \psi' \\ \frac{d^2\tilde{\psi}}{dE^2} &= \left(\frac{40E}{128E^3 - 27} \right) \tilde{\psi} \end{aligned} \tag{1.33}$$

Proof: The first two statements are obvious. To prove the last assertion we use the following definitions of other elliptic integrals.

$$\begin{aligned} \tilde{\omega} &:= \oint_C dw \sqrt{w^4 + 2w + 2E} \\ \omega &:= \oint_C dw \frac{1}{\sqrt{w^4 + 2w + 2E}} \\ \omega' &:= - \oint_C dw \frac{1}{w^4 + 2w + 2E^{3/2}} \\ \tilde{\phi} &:= \oint_C dw w \sqrt{w^4 + 2w + 2E} \end{aligned}$$

$$\begin{aligned}\phi &:= \oint_C dw \frac{w}{\sqrt{w^4 + 2w + 2E}} \\ \phi' &:= - \oint_C dw \frac{w}{w^4 + 2w + 2E^{3/2}}\end{aligned}$$

That these are interrelated can be shown by integration by parts as follows:

$$\begin{aligned}\tilde{\psi} &= -\frac{1}{3} \oint_C dw w^3 \frac{2w^3 + 1}{\sqrt{P(w)}} \\ &= -\frac{2}{3} \oint_C dw w^2 \frac{w^4 + 2w + 2E - (3/2)w - 2E}{\sqrt{P(w)}} \\ &= -\frac{2}{3} \tilde{\psi} + \oint_C dw \frac{w^3}{\sqrt{P(w)}} + \frac{4E}{3} \psi \\ &= -\frac{2}{3} \tilde{\psi} + \frac{1}{2} \oint_C dw \frac{2w^3 + 1}{\sqrt{P(w)}} - \frac{1}{2} \omega + \frac{4E}{3} \psi\end{aligned}$$

Therefore, we get

$$\omega = \frac{8}{3} E \psi - \frac{10}{3} \tilde{\psi}. \quad (1.34)$$

Similarly, starting with $\tilde{\omega}$ and integrating by parts gives

$$3\tilde{\omega} = 3\phi + 4E\omega, \quad (1.35)$$

and starting with $\tilde{\phi}$ gives

$$2\tilde{\phi} = \frac{3}{2} \psi + 2E\phi. \quad (1.36)$$

Finally, starting with ϕ , we get

$$\phi' = -\frac{4}{3} E \phi'. \quad (1.37)$$

Differentiating Eqn(1.34) w.r.t. E yields

$$\omega' = \frac{8}{3} E \psi' - \frac{2}{3} \psi. \quad (1.38)$$

Doing the same with Eqn(1.35) gives

$$-\omega = 3\phi' + 4E\omega'. \quad (1.39)$$

Now eliminating $\tilde{\omega}$, ω , ω' between Eqns(1.34-1.39) gives the desired differential equation for $\psi(E)$. $\square$

Note that these show that $\tilde{\omega}_j$ and, therefore, the periods ω_j are singular at the zeroes of $128E^3 - 27$. These are precisely the degenerate values D_k found in Eqn(2.12).

Consider $\tilde{\psi}_j$, in the degenerate limit $E \rightarrow D_k$ (for some k). We drop the subscript j below, and consider momentarily only the singular case. Fröbenius expansion of the solutions $\tilde{\psi}$ of Eqn(1.33) show that there exists a singular solution with behaviour

$$\tilde{\psi} \approx c_0 + c_1(E - D_k) \ln(E - D_k)$$

as $E \rightarrow D_k$, where c_0 is an arbitrary constant and c_1 is a multiple of it. (For a derivation of similar results with more detail see Appendix B of [5].) Moreover, we have

$$\psi = \tilde{\psi}'(E) \sim c_1 \ln(E - D_k)$$

as $E \rightarrow D_k$. The period in this case is given by

$$\omega = \frac{8}{3}E\psi - \frac{10}{3}\tilde{\psi} \approx \frac{8D_k c_1}{3} \ln(E - D_k). \quad (1.40)$$

There is also a linearly dependent solution analytic at D_k with expansion

$$\omega_j \approx b_0 + b_1(E - D_k)$$

as $E \rightarrow D_k$, where b_0 is an arbitrary constant and b_1 is a multiple of it.

References

- [1] E. Ince. *Ordinary Differential Equations*. Dover, New York, 1954.
- [2] P. Boutroux. Recherches sur les transcendentes de M. Painlevé et l'étude asymptotique des équations différentielles du second ordre. *Ann. École Norm.*, 30:265–375, 1913.
- [3] P. Boutroux. Recherches sur les transcendentes de M. Painlevé et l'étude asymptotique des équations différentielles du second ordre. *Ann. École Norm.*, 31:99–159, 1914.

- [4] A. V. Kitaev. Elliptic asymptotics of the first and the second Painlevé transcendents. *Russian Math. Surveys*, 49:81–150, 1994.
- [5] N. Joshi and M. D. Kruskal. The Painlevé connection problem: an asymptotic approach I. *Stud. Appl. Math.*, 86:315–376, 1992.
- [6] B. McCoy. Spin systems, statistical mechanics, and Painlevé functions. In D. Levi and P. Winternitz, editors, *Painlevé transcendents*, NATO ASI Series B, pages 377–391. Plenum Press, 1992.
- [7] M. J. Ablowitz and H. Segur. *Solitons and the Inverse Scattering Transform*. SIAM, Philadelphia, 1981.
- [8] M. J. Ablowitz and P. A. Clarkson. *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, volume 149 of *London Mathematical Society Lecture Notes in Mathematics*. Cambridge University Press, Cambridge, 1991.
- [9] M. J. Ablowitz and H. Segur. Asymptotic solutions of the Korteweg-deVries equation. *Stud. Appl. Math.*, 57:13–44, 1977.
- [10] H. Segur and M. J. Ablowitz. Connection results for the second Painlevé equation. *Physica*, 3D:165–184, 1981.
- [11] N. Joshi and M. D. Kruskal. An asymptotic approach to the connection problem for the first and the second Painlevé equations. *Phys. Lett. A*, 130:129–137, 1988.
- [12] P. A. Clarkson and J. B. McLeod. Integral equations and connection formulæ for the Painlevé equations. In D. Levi and P. Winternitz, editors, *Painlevé transcendents*, NATO ASI Series B, pages 1–31. Plenum Press, 1992.
- [13] A. R. Its, A. S. Fokas, and A.A. Kapaev. On the asymptotic analysis of the Painlevé equations via the isomonodromy method. *Nonlinearity*, 7:1291–1325, 1994.
- [14] P. Deift and X. Zhou. Asymptotics of the Painlevé II equation. *Comm. Pure Appl. Math.*, 48:277–337, 1995.

- [15] A. Bassom, P. A. Clarkson, C. Law, and J. B. McLeod. Applications of uniform asymptotics to the second Painlevé transcendent. 1996. preprint 9609005 in solv-int archive, at URL <http://xyz.lanl.gov/>.
- [16] T. Kawai and Y. Takei. WKB analysis of Painlevé transcendents with a large parameter I. *Adv. Math.*, 118:1–33, 1996.
- [17] T. Kawai and Y. Takei. On the structure of Painlevé transcendents with a large parameter II. *Proc. Japan Acad. Ser. A Math. Sci.*, 72:144–147, 1996.
- [18] T. Kawai and Y. Takei. On the structure of Painlevé transcendents with a large parameter. *Proc. Japan Acad. Ser. A Math. Sci.*, 69:224–229, 1993.
- [19] K. Okamoto. Isomonodromic deformation and Painlevé equations and the garnier system. *J. Fac. Sci. Univ. Tokyo Sec. IA Math*, 33:575–618, 1986.
- [20] R. Fuchs. *Math. Annalen*, 63:301–321, 1907.
- [21] A. V. Kitaev. Turning points of linear systems and double asymptotics of the Painlevé transcendents. In D. Levi and P. Winternitz, editors, *Painlevé transcendents*, NATO ASI Series B, pages 81–96. Plenum Press, 1992.
- [22] N. Joshi and M. D. Kruskal. A direct proof that the solutions of the six Painlevé equations have no movable singularities except poles. *Stud. Appl. Math.*, 93:187–207, 1994.
- [23] P. Painlevé. Mémoire sur les équations différentielles dont l’intégrale générale est uniforme. *Bull. Soc. Math.*, 28:201–261, 1902.
- [24] C. Bender and S. Orszag. *Advanced Mathematical Methods for Scientists and Engineers*. McGraw-Hill, New York, 1978.
- [25] E. Hille. *Ordinary Differential Equations in the Complex Domain*. John Wiley and Sons,, New York, 1976.
- [26] N. Joshi. Asymptotic studies of the Painlevé equations. 1997. to appear in Proceedings of *the Painlevé property, one century*

later, Institute of Geometry and Applications preprint 97:06 at
<http://www.maths.adelaide.edu.au/pure/iga/preprints.html>.